

Irregular semi-convex gradient systems perturbed by noise and application to the stochastic Cahn–Hilliard equation

Giuseppe Da Prato, Scuola Normale Superiore di Pisa,
Piazza dei Cavalieri 7, 56126, Pisa, Italy

Arnaud Debussche, Ecole Normale Supérieure de Cachan,
antenne de Bretagne, Campus de Ker Lann, 35170 Bruz, France.

Luciano Tubaro, Department of Mathematics, University of Trento,
Via Sommarive 14, 38050 Povo, Italy.

Abstract

We prove essential self-adjointness of Kolmogorov operators corresponding to gradient systems with potentials U such that DU is not square integrable with respect to the invariant measure (irregular potentials). An application is given to the Cahn–Hilliard–Cook equation in dimension one. In this case the spectral gap is proved for the corresponding semigroup. We also obtain a log-Sobolev inequality.

1 Introduction and setting of the problem

Let H be a separable real Hilbert space (norm $|\cdot|$, inner product $\langle \cdot, \cdot \rangle$). We are concerned with the following Kolmogorov operator

$$N_0\varphi = \frac{1}{2} \operatorname{Tr} [D^2\varphi(x)] + \langle x, AD\varphi(x) - \langle DU, D\varphi \rangle, \quad \varphi \in \mathcal{E}_A(H),$$

where D denotes the Fréchet derivative with respect to x . Here $A: \mathcal{D}(A) \subset H \rightarrow H$ is a negative self-adjoint operator such that A^{-1} is of trace class and $U: H \rightarrow (-\infty, +\infty]$ is a semi-convex function. Moreover $\mathcal{E}_A(H)$ is the vector space of all linear combinations of functions of the form

$$\cos(\langle x, h \rangle), \sin(\langle x, h \rangle), \quad h \in \mathcal{D}(A).$$

Let μ be the Gaussian measure in H with mean 0 and covariance operator $Q := -\frac{1}{2} A^{-1}$, we consider the measure

$$\nu(dx) = Z^{-1} e^{-2U(x)} \mu(dx), \tag{1.1}$$

where Z is a normalization constant:

$$Z = \int_H e^{-2U(x)} \mu(dx) \quad (1.2)$$

Our goal is to show that, under suitable assumptions on U , the operator N_0 is dissipative in some space $L^p(H, \nu)$, $p \geq 1$ and that its closure is m -dissipative.

As well known, the Kolmogorov operator N_0 is related to a gradient system described by the following differential stochastic equation

$$dX = (AX - DU(X))dt + dW(t), \quad X(0) = x,$$

where $W(t)$ is a cylindrical Wiener process on H .

Several papers have been devoted to gradient systems. We recall the Dirichlet forms approach, [1], [2], [14], and the semigroup approach, see [12] and references therein. But in all these papers the assumption that (at least) DU is square integrable with respect to ν :

$$\int_H |DU(x)|^2 \nu(dx) < +\infty, \quad (1.3)$$

is made. This assumption is fulfilled in several applications as the reaction-diffusion equations, but it does not hold for semilinear equations perturbed by noise where the nonlinearity involves the derivative of the unknown, see [10] for a discussion on this point. In [10] a concrete case, the Kolmogorov equation corresponding to the p -laplacian (perturbed by a bilaplacian) was considered. In the present paper we replace (1.3) with the weaker condition

$$\int_H |(-A)^{-\frac{1}{2+2\beta}} DU|^{2+2\beta} d\nu < +\infty, \quad (1.4)$$

where $0 \leq \beta \leq 1$, proving that the closure $N_{1+\beta}$ of the operator N_0 is m -dissipative in $L^{1+\beta}(H, \nu)$. As an application, we solve the Kolmogorov equation corresponding to the stochastic Cahn-Hilliard equation.

Let us explain our method. Proceeding as in [9], we consider an approximating equation

$$\lambda \varphi_\alpha - \frac{1}{2} \operatorname{Tr} [D^2 \varphi_\alpha(x)] - \langle x, AD \varphi_\alpha(x) \rangle + \langle DU_\alpha, D \varphi_\alpha \rangle = f, \quad (1.5)$$

where $f \in \mathcal{E}_A(H)$, $\lambda > 0$, and U_α is a smooth approximation of U . We prove that $\varphi_\alpha \in D(N_{1+\beta})$, so that it can be written as

$$\lambda\varphi_\alpha - N_{1+\beta}\varphi_\alpha = f + \langle DU - DU_\alpha, D\varphi_\alpha \rangle. \quad (1.6)$$

Now the key point is to show that

$$\lim_{\alpha \rightarrow 0} \langle DU - DU_\alpha, D\varphi_\alpha \rangle = 0 \quad \text{in } L^{1+\beta}(H, \nu), \quad (1.7)$$

so that the range of $\lambda - N_{1+\beta}$ is dense in $L^{1+\beta}(H, \nu)$ and $N_{1+\beta}$ is m -dissipative. In [9], (1.7) was proved using (1.3) and the basic inequality

$$\int_H N_0 \varphi \varphi d\nu = -\frac{1}{2} \int_H |D\varphi|^2 d\nu, \quad \varphi \in \mathcal{E}_A(H), \quad (1.8)$$

which yields easily an a-priori estimate for $\int_H |D\varphi|^2 d\nu$. In the present situation, since only (1.4) holds, (1.8) is no longer sufficient. We need a stronger estimate which is proved in §3.

§2 is devoted to some preliminaries, §4 to an application to the stochastic Cahn–Hilliard equation in the interval $[0, \pi]$. In this case we prove that N_0 is essentially self-adjoint in $L^2(H, \nu)$. Moreover, we prove the Poincaré and the log-Sobolev inequalities for the measure ν . This implies that the spectral gap property holds for N_2 .

We notice that the Poincaré and the log-Sobolev inequalities do not follow from the Bakry–Emery criterion, see [3], due to the lack of regularity of the potential U of the Cahn–Hilliard equation. The main idea to prove these inequalities is to show that ν is the image of a measure ν_0 through the embedding $L^2([0, \pi]) \subset H^{-1}([0, \pi])$ where ν_0 is the invariant measure for a reaction–diffusion system for which the Poincaré and the log-Sobolev inequalities hold.

2 Preliminaries

Let us state our assumptions. Concerning A we shall assume that

Hypothesis 2.1

(i) A is self-adjoint and there exists $\omega > 0$ such that

$$\langle Ax, x \rangle \leq -\omega |x|^2, \quad x \in \mathcal{D}(A).$$

(ii) A^{-1} is of trace class.

Remark. From (ii) it follows that there exist a complete orthonormal system $\{e_k\}$ in H and a sequence of positive numbers $\{\alpha_k\}$ such that

$$Ae_k = -\alpha_k e_k, \quad k \in \mathbb{N}, \quad \text{with} \quad \sum_{k \in \mathbb{N}} \frac{1}{\alpha_k} < +\infty.$$

We consider the operator N_0 as a perturbation of the Ornstein–Uhlenbeck operator L

$$L\varphi(x) = \frac{1}{2} \operatorname{Tr} [D^2\varphi(x)] + \langle x, AD\varphi(x) \rangle, \quad x \in H, \quad \varphi \in \mathcal{E}_A(H),$$

that is

$$N_0\varphi = L\varphi(x) - \langle DU, D\varphi \rangle, \quad \varphi \in \mathcal{E}_A(H).$$

We recall that L is a self-adjoint operator in $L^2(H, \mu)$ with the property that

$$\int_H L\varphi \psi \, d\mu = -\frac{1}{2} \int_H \langle D\varphi, D\psi \rangle \, d\mu, \quad \varphi, \psi \in W^{1,2}(H, \mu). \quad (2.1)$$

Concerning U we shall make two assumptions.

Hypothesis 2.2

- (i) Given $U: H \rightarrow (-\infty, +\infty]$, there exists $\delta > 0$ such that the function $x \rightarrow U(x) + \delta|x|^2$ is convex.
- (ii) The number Z defined by (1.2) is finite and positive.
- (iii) There exists a family $\{U_\alpha\}_{\alpha>0}$ of C^2 class functions such that $x \rightarrow U_\alpha(x) + \delta|x|^2$ is convex, $U_\alpha(x) \leq U(x)$ and $U_\alpha(x) \uparrow U(x)$ for any $x \in H$.

We shall denote by ν_α the Borel measure in H defined as

$$\nu_\alpha(dx) = Z_\alpha^{-1} e^{-2U_\alpha(x)} \mu(dx),$$

where

$$Z_\alpha := \int_H e^{-2U_\alpha(x)} \mu(dx).$$

Hypothesis 2.3

- (i) $\lim_{\alpha \rightarrow 0} (-A)^{-\frac{1}{2+2\beta}} DU_\alpha =: (-A)^{-\frac{1}{2+2\beta}} DU$ in $L^{2+2\beta}(H, \nu; H)$.
- (ii) $\lim_{\alpha \rightarrow 0} \int_H |(-A)^{-\frac{1}{2+2\beta}} DU_\alpha - (-A)^{-\frac{1}{2+2\beta}} DU|^{2+2\beta} d\nu_\alpha = 0$.
- (iii) If $\beta = 0$, we also assume that there exists $\varepsilon > 0$ such that $(-A)^{-\frac{1}{2}} DU \in L^{2+\varepsilon}(H, \nu; H)$.

We set

$$\mathcal{N}_\alpha \varphi = L\varphi - \langle DU_\alpha, D\varphi \rangle, \quad \varphi \in \mathcal{E}_A(H).$$

Lemma 2.4 *The following identity holds*

$$\int_H \mathcal{N}_\alpha \varphi \psi d\nu_\alpha = -\frac{1}{2} \int_H \langle D\varphi, D\psi \rangle d\nu_\alpha, \quad \varphi, \psi \in \mathcal{E}_A(H). \quad (2.2)$$

In particular, taking $\psi = 1$, we get

$$\int_H \mathcal{N}_\alpha \varphi d\nu_\alpha = 0, \quad \varphi \in \mathcal{E}_A(H). \quad (2.3)$$

that is ν_α is infinitesimally invariant for \mathcal{N}_α .

Proof. Let $\varphi, \psi \in \mathcal{E}_A(H)$. We have by (2.1)

$$\begin{aligned} \int_H L\varphi \psi d\nu_\alpha &= \int_H L\varphi(\psi \rho_\alpha) d\mu = -\frac{1}{2} \int_H \langle D\varphi, D(\psi \rho_\alpha) \rangle d\mu \\ &= -\frac{1}{2} \int_H \langle D\varphi, D\psi \rangle d\nu_\alpha + \int_H \langle D\varphi, DU_\alpha \rangle \psi d\nu_\alpha = 0, \end{aligned}$$

and the conclusion follows. ■

We can now prove that the measure ν is infinitesimally invariant for N_0 .

Proposition 2.5 *We have*

$$\int_H N_0 \varphi \, d\nu = 0, \quad \varphi \in \mathcal{E}_A(H), \quad (2.4)$$

and

$$\int_H N_0 \varphi \, \varphi \, d\nu = -\frac{1}{2} \int_H |D\varphi|^2 d\nu, \quad \varphi \in \mathcal{E}_A(H). \quad (2.5)$$

Proof. It is enough to prove (2.4), (2.5) follows if we take φ^2 in (2.4). But this follows from (2.3) letting α tend to 0 and taking into account Hypothesis 2.3–(ii). ■

Proposition 2.6 *N_0 is dissipative in $L^{1+\beta}(H, \nu)$.*

Proof. The proof is standard, see [13]. ■

3 m –dissipativity of $N_{1+\beta}$

Let us first note that, thanks to Proposition 2.6, N_0 is closable in $L^{1+\beta}(H, \nu)$; we denote by $N_{1+\beta}$ its closure. We are going to show that $N_{1+\beta}$ is m –dissipative.

Let $\alpha, \lambda > 0$, $f \in C_b^1(H)$ and consider the approximating equation

$$\lambda \varphi_\alpha - L_{1+\beta} \varphi_\alpha + \langle DU_\alpha, D\varphi_\alpha \rangle = f, \quad \lambda > 0. \quad (3.1)$$

Lemma 3.1 *If $\lambda > \delta$ equation (3.1) has a unique solution $\varphi_\alpha \in C_b^1(H) \cap \mathcal{D}(N_{1+\beta})$ and*

$$N_{1+\beta} \varphi_\alpha(x) = L_{1+\beta} \varphi_\alpha(x) - \langle DU(x), D\varphi_\alpha(x) \rangle, \quad x \in H, \quad (3.2)$$

and

$$\|D\varphi_\alpha\|_0 \leq \frac{1}{\lambda - \delta} \|Df\|_0, \quad (3.3)$$

where $\|\cdot\|_0$ denotes the sup norm.

Proof. **Step 1.** $\varphi_\alpha \in C_b^1(H)$ and (3.3) holds.

It is well known that

$$\varphi_\alpha(x) = \int_0^\infty e^{-\lambda t} \mathbb{E}[f(X_\alpha(t, x))] dt$$

where $X_\alpha(t, x)$ is the solution to the following stochastic differential equation

$$dX_\alpha = (AX_\alpha - DU_\alpha(X_\alpha))dt + dW_t, \quad X_\alpha(0) = x.$$

Then for any $h \in H$

$$\langle D\varphi_\alpha(x), h \rangle = \int_0^\infty e^{-\lambda t} \mathbb{E}[\langle Df(X_\alpha(t, x)), \eta_\alpha^h(t, x) \rangle] dt \quad (3.4)$$

where $\eta_\alpha^h(t, x)$ is the solution to the following equation

$$\frac{d}{dt} \eta_\alpha^h = A\eta_\alpha^h - D^2U_\alpha(X_\alpha) \cdot \eta_\alpha^h, \quad \eta_\alpha^h(0) = h. \quad (3.5)$$

Consequently $\varphi_\alpha \in C_b^1(H)$. Moreover, multiplying both sides of equation (3.5) by η_α^h and taking in account the dissipativity of A and the convexity of $x \rightarrow U_\alpha(x) + \delta|x|^2$, yields

$$|\eta_\alpha^h| \leq e^{\delta t} |h|, \quad t > 0.$$

Using (3.4) we get

$$|\langle D\varphi_\alpha(x), h \rangle| \leq \frac{1}{\lambda - \delta} |h| \|Df\|_0,$$

and (3.3) is proved.

Step 2. $\varphi_\alpha \in \mathcal{D}(L_{1+\beta})$ where $L_{1+\beta}$ is the infinitesimal generator of the Ornstein–Uhlenbeck semigroup in $L^{1+\beta}(H, \mu)$,

$$R_t\varphi(x) = \int_H \varphi(e^{tA}x + y) N(0, Q_t)(dy), \quad \varphi \in C_b(H),$$

where

$$Q_t = -\frac{1}{2}A^{-1}(1 - e^{2tA}), \quad t \in [0, +\infty];$$

and observing that $Q_\infty = Q$.

We need a further approximating equation:

$$\lambda \varphi_{\alpha,\beta} - L_{1+\beta} \varphi_{\alpha,\beta} + \frac{1}{1+\beta|DU_\alpha|^2} \langle DU_\alpha, D\varphi_{\alpha,\beta} \rangle = f, \quad \lambda > 0. \quad (3.6)$$

By [12, Proposition 6.6.4], equation (3.6) has a unique solution $\varphi_{\alpha,\beta} \in C_b(H)$. Moreover

$$\|\varphi_{\alpha,\beta}\|_0 \leq \|f\|_0,$$

and there exists $C > 0$ such that

$$\|D\varphi_{\alpha,\beta}\|_0 \leq \frac{C}{\lambda - \delta} \|f\|_1.$$

Since DU_α has linear growth, there exists $C(\alpha, \|f\|_1) > 0$ such that

$$|L_{1+\beta} \varphi_{\alpha,\beta}(x)| \leq C(\alpha, \|f\|_1)(1 + |x|), \quad x \in H.$$

It follows that

$$\int_H |L_{1+\beta} \varphi_{\alpha,\beta}(x)|^2 \mu(dx) \leq C(\alpha, \|f\|_1)(1 + \text{Tr } Q).$$

By a standard argument this implies that $\varphi_\alpha \in \mathcal{D}(L_{1+\beta})$.

Step 3. $\varphi_\alpha \in \mathcal{D}(N_{1+\beta})$ and (3.2) holds.

Let us first consider the case when $\beta \in (0, 1]$. We recall that, see [11],

$$\langle DR_t \varphi, h \rangle = \int_H \langle \Lambda(t)h, Q_t^{-1/2}y \rangle \varphi(e^{tA}x + y) N(0, Q_t)(dy), \quad \varphi \in L^p(H, \mu),$$

with

$$\Lambda(t) = Q_t^{-1/2} e^{tA}.$$

Hence,

$$\langle (-A)^{\frac{1}{2+2\beta}} DR_t \varphi, h \rangle = \int_H \langle (-A)^{\frac{1}{2+2\beta}} \Lambda(t)h, Q_t^{-1/2}y \rangle \varphi(e^{tA}x + y) N(0, Q_t)(dy)$$

and, for $p \geq 1$,

$$\begin{aligned} |\langle (-A)^{\frac{1}{2+2\beta}} DR_t \varphi, h \rangle|^p &\leq \int_H |\varphi(e^{tA}x + y)|^p N(0, Q_t)(dy) \\ &\quad \left(\int_H |\langle (-A)^{\frac{1}{2+2\beta}} \Lambda(t)h, Q_t^{-1/2}y \rangle|^q N(0, Q_t)(dy) \right)^{\frac{p}{q}}, \end{aligned}$$

where q is the conjugate exponent of p . It follows that

$$|(-A)^{\frac{1}{2+2\beta}} D R_t \varphi|^p \leq C t^{-p \frac{2+\beta}{2+2\beta}} R_t(\varphi)^p(x), \quad \varphi \in L^p(H, \mu),$$

which, integrating with respect to μ and taking the Laplace tranform, yields

$$\|(-A)^{\frac{1}{2+2\beta}} D(\lambda - L^{1+\beta})f\|_{L^p(H, \mu)} \leq C(\lambda) \|f\|_{L^p(H, \mu)}, \quad \varphi \in L^p(H, \mu). \quad (3.7)$$

We are now ready to prove that that $\varphi_\alpha \in \mathcal{D}(N_{1+\beta})$.

Since $\mathcal{E}_A(H)$ is a core for $L_{1+\beta}$, see [5], there exists a sequence $\{\varphi_n\} \subset \mathcal{E}_A(H)$ such that,

$$\lim_{n \rightarrow \infty} \varphi_n = \varphi_\alpha, \quad \lim_{n \rightarrow \infty} L_{1+\beta} \varphi_n = L_{1+\beta} \varphi_\alpha, \quad \text{in } L^{1+\beta}(H, \mu).$$

We claim that

$$\lim_{n \rightarrow \infty} N_{1+\beta} \varphi_n = L_{1+\beta} \varphi_\alpha - \langle DU(x), D\varphi_\alpha \rangle \quad \text{in } L^{1+\beta}(H, \nu),$$

which proves that $\varphi_\alpha \in \mathcal{D}(N_{1+\beta})$.

Since by (3.7) we know that $(-A)^{\frac{1}{2+2\beta}} D\varphi_n \rightarrow (-A)^{\frac{1}{2+2\beta}} D\varphi$ in $L^{1+\beta}(H, \mu)$, it is enough to show , in view of the Vitali theorem, that

$$\int_H |\langle DU(x), D\varphi_n \rangle|^{1+\beta+\varepsilon} \nu(dx),$$

is bounded, for some $\varepsilon > 0$. We have in fact

$$\begin{aligned} & \int_H |\langle DU(x), D\varphi_n \rangle|^{1+\beta+\varepsilon} \nu(dx) \\ & \leq \int_H |(-A)^{-\frac{1}{2+2\beta}} DU(x)|^{1+\beta+\varepsilon} |(-A)^{\frac{1}{2+2\beta}} D\varphi_n|^{1+\beta+\varepsilon} \nu(dx) \\ & \leq \left(\int_H |(-A)^{-\frac{1}{2+2\beta}} DU(x)|^{2+2\beta} \nu(dx) \right)^{\frac{1+\beta+\varepsilon}{2+2\beta}} \\ & \quad \times \left(\int_H |(-A)^{\frac{1}{2+2\beta}} D\varphi_n|^{\frac{2(1+\beta+\varepsilon)(1+\beta)}{1+\beta-\varepsilon}} \nu(dx) \right)^{\frac{1+\beta-\varepsilon}{2+2\beta}}. \end{aligned}$$

Now the claim follows from (3.7).

Let us now consider the case $\beta = 0$.

Since $\mathcal{E}_A(H)$ is a core for L_2 , there exists a sequence $\{\varphi_n\} \subset \mathcal{E}_A(H)$ such that,

$$\lim_{n \rightarrow \infty} \varphi_n = \varphi_\alpha, \quad \lim_{n \rightarrow \infty} L_2 \varphi_n = L_2 \varphi_\alpha, \quad \text{in } L^2(H, \mu).$$

It follows that for all $\psi \in \mathcal{D}(L_2)$ we have, see [12, page 215], $|(-A)^{1/2} D\psi| \in L^2(H, \mu)$ and there exists $c > 0$ such that

$$\int_H |(-A)^{1/2} D\psi|^2 d\mu \leq c \int_H |L_2 \psi|^2 d\mu, \quad \forall \psi \in \mathcal{D}(L_2). \quad (3.8)$$

Consequently, by (3.8) it follows that

$$\lim_{n \rightarrow \infty} (-A)^{1/2} D\varphi_n = (-A)^{1/2} D\varphi_\alpha \quad \text{in } L^2(H, \mu; H), \quad (3.9)$$

and there exists $c > 0$ such that

$$\int_H |(-A)^{1/2} D\varphi_n|^2 d\mu \leq c.$$

We claim that

$$\lim_{n \rightarrow \infty} N_0 \varphi_n = L_2 \varphi_\alpha - \langle DU(x), D\varphi_\alpha \rangle \quad \text{in } L^1(H, \nu).$$

This will imply that $\varphi_\alpha \in \mathcal{D}(N_1)$. It is enough to show that

$$\int_H |\langle DU(x), D\varphi_n(x) \rangle|^{1+\gamma} d\nu,$$

is bounded, for some $\gamma > 0$. We have in fact

$$\begin{aligned} & \int_H |\langle DU(x), D\varphi_n(x) \rangle|^{1+\gamma} d\nu \leq \\ & \left(\int_H |(-A)^{-\frac{1}{2}} DU(x)|^{2+\frac{4\gamma}{1-\gamma}} d\nu \right)^{\frac{1-\gamma}{2}} \left(\int_H |(-A)^{\frac{1}{2}} D\varphi_n|^2 d\nu \right)^{\frac{1+\gamma}{2}}, \end{aligned}$$

hence, because of Hypothesis (2.3)-(iii) in the case $\beta = 0$, we can apply the Vitali theorem choosing $\gamma = \frac{\varepsilon}{4+\varepsilon}$. ■

The following identity for $D\varphi_\alpha$ is central in the proof of our main result.

Proposition 3.2 *Let $f \in C_b^2(H)$ and let φ_α be the solution of (3.1). Then we have*

$$\begin{aligned} & \lambda \int_H |D\varphi_\alpha|^2 d\nu_\alpha + \frac{1}{2} \int_H \text{Tr} [(D^2\varphi_\alpha)^2] d\nu_\alpha + \\ & \int_H |(-A)^{1/2} D\varphi_\alpha|^2 d\nu_\alpha + \int_H \langle D^2 U_\alpha D\varphi_\alpha, D\varphi_\alpha \rangle d\nu_\alpha = \\ & \int_H \langle D\varphi_\alpha, Df \rangle d\nu_\alpha = 2 \int_H f(f - \lambda\varphi_\alpha) d\nu_\alpha. \end{aligned} \quad (3.10)$$

Proof. Let $f \in C_b^1(H)$ and let φ_α be the solution of (3.1). Let us differentiate both sides of (3.1) with respect to D_k , $k \in \mathbb{N}$, where D_k is the derivative in the direction of e_k . We obtain

$$\lambda D_k \varphi_\alpha - L D_k \varphi_\alpha + \langle D U_\alpha, D D_k \varphi_\alpha \rangle + \mu_k D_k \varphi_\alpha + \langle D D_k U_\alpha, D \varphi_\alpha \rangle = D_k f.$$

Multiplying by $D_k \varphi_\alpha$, integrating with respect to ν_α and taking into account (2.2), we find that

$$\begin{aligned} & \lambda \int_H |D_k \varphi_\alpha|^2 d\nu_\alpha + \frac{1}{2} \int_H |D D_k \varphi_\alpha|^2 d\nu_\alpha \\ & + \mu_k \int_H |D_k \varphi_\alpha|^2 d\nu_\alpha + \int_H \langle D D_k U_\alpha, D \varphi_\alpha \rangle D_k \varphi_\alpha d\nu_\alpha \\ & = \int_H D_k \varphi_\alpha D_k f d\nu_\alpha. \end{aligned}$$

Summing up on k gives, taking again into account (2.2), the conclusion follows. ■

Corollary 3.3 *There exists $c_1 > 0$ such that for any $f \in C_b^2(H)$*

$$\int_H |(-A)^{1/2} D\varphi_\alpha|^2 d\nu_\alpha \leq c_1 \|f\|_0^2,$$

where φ_α is the solution to (3.1).

Theorem 3.4 *The closure $N_{1+\beta}$ of N_0 in $L^{1+\beta}(H, \nu)$, is m -dissipative in $L^{1+\beta}(H, \nu)$.*

Proof. Let $\lambda > \delta$, $f \in C_b^2(H)$, $\alpha > 0$, and let φ_α be the solution to (3.1). Since by Lemma 3.1 $\varphi_\alpha \in \mathcal{D}(N_{1+\beta})$ we can write

$$\lambda \varphi_\alpha - N_{1+\beta} \varphi_\alpha = \langle (-A)^{-1/2} (DU - DU_\alpha), (-A)^{1/2} D\varphi_\alpha \rangle + f.$$

We claim that

$$\lim_{\alpha \rightarrow 0} \langle DU - DU_\alpha, D\varphi_\alpha \rangle = 0 \quad \text{in } L^{1+\beta}(H, \nu), \quad (3.11)$$

This will conclude the proof by applying the classical result of Lumer and Phillips, [15].

Let us prove (3.11). Since $U_\alpha(x) \leq U(x)$ and $\lim_{\alpha \rightarrow 0} Z_\alpha = Z$, Corollary 3.3 implies

$$\begin{aligned} \int_H |(-A)^{1/2} D\varphi_\alpha|^2 d\nu &= Z^{-1} \int_H |(-A)^{1/2} D\varphi_\alpha|^2 e^{-2U(x)} \mu(dx) \\ &\leq Z^{-1} \int_H |(-A)^{1/2} D\varphi_\alpha|^2 e^{-2U_\alpha(x)} \mu(dx) \leq \frac{Z_\alpha}{Z} c_1 \leq c, \end{aligned}$$

where c is a suitable positive constant. Then, by the Hölder inequality we obtain,

$$\begin{aligned} &\int_H |\langle DU - DU_\alpha, D\varphi_\alpha \rangle|^{1+\beta} d\nu \\ &\leq \left[\int_H |((-A)^{-\frac{1}{2+2\beta}} (DU - DU_\alpha))|^{2+2\beta} d\nu \right]^{\frac{1}{2}} \left[\int_H |(-A)^{\frac{1}{2+2\beta}} D\varphi_\alpha|^{2+2\beta} d\nu \right]^{\frac{1}{2}}. \end{aligned}$$

Now we use the well known interpolatory estimate

$$|(-A)^{\frac{1}{2+2\beta}} x|^{2+2\beta} \leq C |x|^{2\beta} |(-A)^{\frac{1}{2}} x|^2, \quad x \in \mathcal{D}((-A)^{\frac{1}{2}}),$$

and find

$$\begin{aligned}
& \int_H |\langle DU - DU_\alpha, D\varphi_\alpha \rangle|^{1+\beta} d\nu \\
& \leq C \left[\int_H |((-A)^{-\frac{1}{2+2\beta}}(DU - DU_\alpha))|^{2+2\beta} d\nu \right]^{\frac{1}{2}} \left[\int_H |D\varphi_\alpha|^{2\beta} |(-A)^{\frac{1}{2}} D\varphi_\alpha|^2 d\nu \right]^{\frac{1}{2}} \\
& \leq \frac{C}{(\lambda - \delta)^\beta} \|Df\|_0^\beta \|f\|_0 \left(\int_H |(-A)^{-\frac{1}{2+2\beta}}(DU - DU_\alpha)|^{2+2\beta} d\nu \right)^{1/2}.
\end{aligned}$$

thanks to (3.3) and Corollary 3.3. The proof is complete thanks to Hypothesis 2.3 *i*). ■

4 The stochastic Cahn–Hilliard equation

4.1 m –dissipativity

The Cahn–Hilliard equation is a phenomenological model for various types of non equilibrium phase transitions as the early stage of *spinodal decomposition*, a physical phenomenon that arises when we rapidly quench an alloy from the stable region (high temperature) to the unstable region (low temperature). Cook took into account also the thermal fluctuations introducing the stochastic Cahn–Hilliard equation, which in the litterature is known also as the Cahn–Hilliard–Cook equation.

This equation has been intensively studied, see e.g. [4], [6], [7], and the references cited therein.

We will apply the abstract results of §3 to the following stochastic Cahn–Hilliard equation:

$$\left\{ \begin{array}{l} dX = D_\xi^2(-D_\xi^2 X + f(X))dt + dW(t), \quad \text{in } [0, +\infty) \times [0, \pi], \\ \int_0^\pi X(\xi) d\xi = 0 \\ D_\xi X(t, 0) = D_\xi^3 X(t, 0) = D_\xi X(t, \pi) = D_\xi^3 X(t, \pi) = 0, \\ X(0, \cdot) = x, \end{array} \right. \quad (4.1)$$

where $W(t)$ is a cylindrical white noise (in time and in space), and where $f \in C^2(\mathbb{R})$ is such that

$$|f(r)| \leq a(1 + |r|^{2m-1}), \quad (4.2)$$

for some a and the function

$$g(r) = \int_0^r f(s)ds$$

is semiconvex. Typically g is a polynomial with positive leading coefficient of even order (greater than or equal to 4). In order to avoid technical complications below, we make the additional assumption that f is monotone, however all our results hold in the more general case of the derivative of a semiconvex function.

The stochastic Cahn–Hilliard equation with periodic boundary conditions can be treated in the same way.

It would be interesting to study the Cahn–Hilliard equation in two (or more) dimensions. This would require a renormalization procedure, which we will try to study in a future paper.

In general X denotes concentration, for instance in the case of a binary alloy (Cu,Zn), X can be the concentration of Cu. In the deterministic case the Cahn–Hilliard equation has the property that the total concentration - which corresponds to the spatial average of X - is a conserved quantity. Without loss of generality, we assume that this average is zero. It is natural to require that the noise does not destroy this property. Thus we work in spaces of zero average functions and introduce $\mathcal{H}^1(0, \pi)$, the space of functions in $H^1(0, \pi)$ whose average is zero, and its dual $\mathcal{H}^{-1}(0, \pi)$.

It is natural to study this problem in the space $H = \mathcal{H}^{-1}(0, \pi)$ because, with this choice, the equation is of gradient type and the corresponding transition semigroup is reversible.

We also consider the Hilbert space $\dot{L}^2(0, \pi)$ of all square integrable functions φ on $[0, \pi]$ with zero average. Its inner product is denoted by $\langle \cdot, \cdot \rangle$.

Let $\{e_k\}_{k \in \mathbb{N}^*}$ (¹) be the orthonormal basis on $\dot{L}^2(0, \pi)$ defined by

$$e_k(\xi) = (\pi/2)^{-1/2} \cos(k\xi), \quad k \in \mathbb{N}^*,$$

and, for any $x \in L^2(0, \pi)$, set

$$x_k = \langle x, e_k \rangle, \quad k \in \mathbb{N}^*.$$

¹ $\mathbb{N}^* = 1, 2, \dots$

We shall identify $\dot{L}^2(0, \pi)$ with $\ell^2(\mathbb{N}^*)$ and then we shall consider $\dot{L}^2(0, \pi)$ as a subspace of $\mathbb{R}^{\mathbb{N}^*}$.

Moreover, for any $r \in \mathbb{R}$ we shall denote by \mathcal{H}^r the subspace of $\mathbb{R}^{\mathbb{N}^*}$ of all sequences $x = \{x_k\}_{k \in \mathbb{N}^*}$ such that

$$|x|_r^2 := \sum_{k \in \mathbb{N}^*} (1 + |k|^2)^r |x_k|^2 < +\infty.$$

\mathcal{H}^r is a Hilbert space with the inner product

$$\langle x, y \rangle_r := \sum_{k \in \mathbb{N}^*} |k|^{2r} x_k y_k, \quad x, y \in \mathcal{H}^r. \quad (4.3)$$

The corresponding norm is denoted by $|\cdot|_r$. Notice that $\dot{L}^2(0, \pi) = \mathcal{H}^0$, $\mathcal{H}^1(0, \pi) = \mathcal{H}^1$, $\mathcal{H}^{-1}(0, \pi) = \mathcal{H}^{-1}$ and setting

$$f_k(\xi) = (1 + |k|^2)^{1/2} e_k(\xi), \quad k \in \mathbb{N}^*,$$

then $\{f_k\}_{k \in \mathbb{N}^*}$ is a complete orthonormal basis on \mathcal{H}^{-1} . Clearly, if $r_1 \geq r_2$,

$$|x|_{r_1} \leq |x|_{r_2}.$$

Moreover we assume that $W(t)$ is the cylindrical Wiener process on \mathcal{H}^{-1} defined (formally) by

$$W(t) = \sum_{k \in \mathbb{N}^*} f_k \beta_k,$$

where $\{\beta_k\}_{k \in \mathbb{N}^*}$ is a sequence of mutually independent standard Brownian motions.

Let us define the linear (unbounded) operators A and B in $H = \mathcal{H}^{-1}$ by setting

$$Bf_k = k^2 f_k, \quad k \in \mathbb{N}^*,$$

and

$$Af_k = -k^4 f_k, \quad k \in \mathbb{N}^*.$$

Notice that

$$\mathcal{D}(B) = \mathcal{H}^1, \quad \mathcal{D}(A) = \mathcal{H}^3,$$

and that $B = (-A)^{1/2}$.

Moreover, let us introduce the potential $U: \mathcal{H}^{-1} \mapsto [0, +\infty]$

$$U(x) = \begin{cases} \int_0^\pi g(x(\xi)) d\xi, & \text{if } x \in \mathcal{D}(U), \\ +\infty & \text{otherwise,} \end{cases} \quad (4.4)$$

where

$$\mathcal{D}(U) = \{y \in \dot{L}^2([0, \pi]) : g(y) \in L^1([0, \pi])\},$$

and $g(r) = \int_0^r f(s)ds$.

We have

$$DU(x) \cdot y = \int_0^\pi g(\xi)d\xi.$$

We denote by $D_{\mathcal{H}^r}$ the gradient in \mathcal{H}^r , then

$$D_{\mathcal{H}^{-1}}U = (-A)^{1/2}DU = Bf(x).$$

For $r = -1$, we also set $D = D_{\mathcal{H}^{-1}}$. Thus, equation (4.1) can be written as

$$\begin{cases} dX = (AX - DU(X))dt + dW(t), \\ X(0) = x, \end{cases} \quad (4.5)$$

Now we can consider the Kolmogorov operator

$$N_0\varphi(x) = \frac{1}{2} \operatorname{Tr} [D^2\varphi(x)] + \langle x, AD\varphi(x) \rangle - \langle DU(x), D\varphi(x) \rangle, \quad \varphi \in \mathcal{E}_A(H)$$

which we shall write also as

$$N_0\varphi(x) = \frac{1}{2} \operatorname{Tr} [D^2\varphi(x)] + \langle x, AD\varphi(x) \rangle + \langle f(x), BD\varphi(x) \rangle. \quad (4.6)$$

We set $\mu = N_Q$ where $Q = -\frac{1}{2} A^{-1}$. We have

Theorem 4.1 *Let N_0 be the Kolmogorov operator defined by (4.6), and let ν the probability measure defined by (1.1). Then N_0 is essentially self-adjoint in $L^2(\mathcal{H}^{-1}, \nu)$.*

Proof. We shall apply Theorem 3.4, verifying the required assumptions for $\beta = 1$ and $\delta = 0$.

Verification of Hypothesis 2.1. It follows from the identity

$$\operatorname{Tr} Q = \frac{1}{2} \sum_{k \in \mathbb{N}^*} k^{-4} < +\infty.$$

Verification of Hypothesis 2.2–(ii). For this it is convenient to write $x(\xi)$ in a suitable form. Given $x \in \mathcal{H}^{-1}$ we start from the obvious identity

$$x(\xi) = \sum_{k \in \mathbb{N}^*} \langle x, f_k \rangle_{-1} f_k = \frac{1}{\sqrt{2}} \left\langle Q^{-1/2} x, \sum_{k \in \mathbb{N}^*} \frac{1}{k^2} f_k(\xi) f_k \right\rangle_{-1} = \rho(\xi) W_{\eta_\xi}(x),$$

where W represents the white noise function, η_ξ is the element in \mathcal{H}^{-1} defined by

$$\eta_\xi = \frac{1}{\sqrt{2}\rho(\xi)} \sum_{k \in \mathbb{N}^*} \frac{1}{k^2} f_k(\xi) f_k, \quad (4.7)$$

and

$$\rho^2(\xi) = \frac{1}{2} \sum_{k \in \mathbb{N}^*} \frac{1+k^2}{k^4} e_k(\xi)^2. \quad (4.8)$$

Now we can prove that

$$Z = \int_H e^{-2U(x)} \mu(dx) > 0.$$

For this it is enough to show that

$$\int_H U(x) \mu(dx) < +\infty. \quad (4.9)$$

We have in fact

$$\begin{aligned} \int_H U(x) \mu(dx) &= \int_H \int_0^\pi g(x(\xi)) d\xi \mu(dx) = \int_0^\pi d\xi \int_H g(\rho(\xi) W_{\eta_\xi}(x)) \mu(dx) \\ &= (2\pi)^{-\frac{1}{2}} \int_0^\pi d\xi \int_{-\infty}^{+\infty} e^{-\frac{r^2}{2}} g(\rho(\xi)r) dr < +\infty, \end{aligned}$$

in view of (4.2), and Hypothesis 2.2–(ii) is fulfilled.

Verification of Hypothesis 2.2–(iii).

Let us define approximations U_α of U . Let g_α be the Moreau-Yosida approximations of g

$$g_\alpha(r) = \inf \left\{ g(s) + \frac{1}{2\alpha} (r-s)^2 : s \in \mathbb{R} \right\}.$$

We set

$$U_\alpha(x) = \int_0^\pi g_\alpha((1+\alpha B)^{-1} x(\xi)) d\xi, \quad \alpha > 0. \quad (4.10)$$

Then U_α is of class C^2 . Moreover, $U_\alpha \leq U$. In fact, since

$$(1 + \alpha B)^{-1}x(\xi) = \int_0^\pi k(\xi, \eta) x(\eta) d\eta$$

with $k(\xi, \eta) \geq 0$, we have that $\int_0^\pi k(\xi, \eta) d\eta = 1$: this allows us to apply Jensen inequality to get

$$g_\alpha((1 + \alpha B)^{-1}x) \leq g((1 + \alpha B)^{-1}x) \leq (1 + \alpha B)^{-1}g(x).$$

Hence

$$U_\alpha(x) \leq U(x).$$

Verification of Hypothesis 2.2-(iv). Firstly we observe that

$$DU_\alpha(x) \cdot y = \int_0^\pi g'_\alpha((1 + \alpha B)^{-1}x(\xi)) (1 + \alpha B)^{-1}y(\xi) d\xi,$$

so that

$$D_{\mathcal{H}^{-1}}U_\alpha = (-A)^{1/2}D_{\mathcal{H}^0}U_\alpha = B(1 + \alpha B)^{-1}g'_\alpha((1 + \alpha B)^{-1}).$$

We have to show that

$$\lim_{\alpha \rightarrow 0} \int_{\mathcal{H}^{-1}} |(-A)^{1/4}D_{\mathcal{H}^{-1}}(U - U_\alpha)|_{-1}^4 d\nu = \lim_{\alpha \rightarrow 0} \int_{\mathcal{H}^{-1}} |D_{\mathcal{H}^0}(U - U_\alpha)|_0^4 d\nu = 0.$$

In view of the dominated convergence theorem it is enough to show that $|D_{\mathcal{H}^0}U_\alpha|_0^4$ can be estimated, uniformly in α by a ν -integrable function. We have in fact, using the Jensen inequality

$$\begin{aligned} |D_{\mathcal{H}^0}U_\alpha|_0^4 &= \left(\int_0^\pi D_{\mathcal{H}^0}U_\alpha(x)(\xi)^2 d\xi \right)^2 = \left(\int_0^\pi f_\alpha((1 + \alpha B)^{-1}x)(\xi)^2 d\xi \right)^2 \\ &\leq \left(\int_0^\pi f((1 + \alpha B)^{-1}x)(\xi)^2 d\xi \right)^2 \leq \left(\int_0^\pi f(x)(\xi)^2 d\xi \right)^2. \end{aligned}$$

It remains to show that

$$\int_{\mathcal{H}^{-1}} \left(\int_0^\pi f(x)(\xi)^2 d\xi \right)^2 d\nu < +\infty.$$

We have in fact, thanks to (4.2), and proceeding as in the proof of (4.9),

$$\begin{aligned}
& \int_{\mathcal{H}^{-1}} \left(\int_0^\pi f(x(\xi))^2 d\xi \right)^2 d\nu \leq \pi \int_0^\pi d\xi \int_{\mathcal{H}^{-1}} f(x(\xi))^4 d\nu \\
& \leq a\pi \left(\pi + \int_0^\pi d\xi \int_{\mathcal{H}^{-1}} (x(\xi))^{8m-4} d\nu \right) \\
& = a\pi \left(\pi + (2\pi)^{-\frac{1}{2}} \frac{(8m-4)!}{2^{4m-2}(4m-2)!} \int_0^\pi \rho(\xi)^{8m-4} d\xi \right),
\end{aligned}$$

which is finite. The proof is complete. ■

4.2 Spectral gap

We consider here the invariant measure ν of the Cahn-Hilliard-Cook equation (4.5) in \mathcal{H}^{-1} , that is

$$\nu(dx) = Z^{-1} \exp(-U(x)) \mu(dx),$$

where U is defined by (4.4).

We recall that for a sufficiently smooth function x , we have the following relationship between the derivatives in $\mathcal{H}^0 = \dot{L}^2(0, \pi)$ and in \mathcal{H}^{-1} :

$$D_{\mathcal{H}^{-1}}x = BD_{\mathcal{H}^0}x.$$

Let T be the natural imbedding of \mathcal{H}^0 into \mathcal{H}^{-1} . It is easily checked that the adjoint T' of T is given by $T'y = -B^{-1}y$.

Let us consider on \mathcal{H}^0 the Gaussian measure $\mu_0 = N(0, Q_0)$ with $Q_0 = -\frac{1}{2} B^{-1}$ and set

$$\nu_0(dy) = Z_0^{-1} \exp(-U(y)) \mu_0(dy)$$

and

$$Z_0 = \int_H \exp(-U(y)) \mu_0(dy).$$

By the Bakry-Emery criterion, [3], see also [8], it follows that ν_0 is the unique invariant measure of the following stochastic differential equation

$$dX = (BX - f(X))dt + dW_0.$$

We need the following lemma

Lemma 4.2 *The image measure of ν_0 through the natural imbedding $T: \mathcal{H}^0 \rightarrow \mathcal{H}^{-1}$ coincides with ν .*

Proof. We first prove that

$$\mu(\mathcal{H}^0) = 1. \quad (4.11)$$

We have in fact

$$\int_{\mathcal{H}^{-1}} |x|_{\mathcal{H}^0}^2 \mu(dx) = \int_{\mathcal{H}^{-1}} |\sqrt{B}x|_{\mathcal{H}^{-1}}^2 \mu(dx) = \text{Tr}(B^{-1}) < +\infty.$$

To prove the lemma it is enough to show that for any Borel bounded function $\varphi: \mathcal{H}^{-1} \rightarrow \mathbb{R}$ we have

$$\int_{\mathcal{H}^0} \varphi(y) \nu_0(dy) = \int_{\mathcal{H}^{-1}} \varphi(x) \nu(dx). \quad (4.12)$$

To prove (4.12) we consider a sequence $\{P_n\}$ of finite dimensional approximations of the identity in \mathcal{H}^0 and we set $P'_n = TP_n$, $n \in \mathbb{N}^*$. Then by the change of variables formula in finite dimensional spaces, we get

$$\int_{P_n \mathcal{H}^0} \varphi(P_n y) e^{-2U(P_n y)} N(0, P_n Q_0)(dy) = \int_{P'_n \mathcal{H}^{-1}} \varphi(P'_n x) e^{-2U(P'_n x)} N(0, P'_n Q)(dx).$$

Now, letting n tend to infinity and taking into account (4.11), we find (4.12). \blacksquare

Let us prove now the Poincaré inequality for the measure ν .

Theorem 4.3 *For any $\varphi \in C_b^1(\mathcal{H}^{-1})$ we have*

$$\int_{\mathcal{H}^{-1}} |\varphi(x) - \bar{\varphi}(x)|^2 \nu(dx) \leq \frac{1}{2} \int_{\mathcal{H}^{-1}} |D_{\mathcal{H}^{-1}} \varphi(x)|_{\mathcal{H}^{-1}}^2 \nu(dx), \quad (4.13)$$

where

$$\bar{\varphi} = \int_{\mathcal{H}^{-1}} \varphi(x) \nu(dx).$$

Proof. It is well known, see [3], [8], that the Poincaré inequality holds for the measure ν_0 . Therefore, for any $\varphi \in C_b^1(\mathcal{H}^0)$ we have

$$\int_{\mathcal{H}^0} |\varphi(y) - \bar{\bar{\varphi}}(y)|^2 \nu_0(dy) \leq \frac{1}{2} \int_{\mathcal{H}^0} |D_{\mathcal{H}^0} \varphi(x)|_{\mathcal{H}^0}^2 \nu_0(dy), \quad (4.14)$$

where

$$\overline{\overline{\varphi}} = \int_{\mathcal{H}^0} \varphi(y) \nu_0(dy).$$

On the other hand we have, by the change of variables formula, that $\overline{\overline{\varphi}} = \overline{\varphi}$. Consequently

$$\begin{aligned} \frac{1}{2} \int_{\mathcal{H}^{-1}} |D_{\mathcal{H}^{-1}} \varphi(x)|_{\mathcal{H}^{-1}}^2 \nu(dx) &= \frac{1}{2} \int_{H^{-1}} |BD_{\mathcal{H}^0} \varphi(x)|_{\mathcal{H}^{-1}}^2 \nu(dx) \\ &\geq \frac{1}{2} \int_{H^{-1}} |D_{\mathcal{H}^0} \varphi(x)|_{\mathcal{H}^0}^2 \nu(dx) = \frac{1}{2} \int_{\mathcal{H}^0} |D_{\mathcal{H}^0} \varphi(x)|_{\mathcal{H}^0}^2 \nu_0(dx) \\ &\geq \int_{\mathcal{H}^0} |\varphi(x) - \overline{\varphi}(x)|^2 \nu_0(dx) = \int_{\mathcal{H}^{-1}} |\varphi(x) - \overline{\varphi}(x)|^2 \nu(dx), \end{aligned}$$

by the change of variables formula. ■

The spectral gap follows now easily, see [3].

Corollary 4.4 *Let N_2 be the closure of N_0 in $L^2(H, \nu)$ and let $\sigma(N_2)$ be its spectrum. Then we have*

$$\sigma(N_2) \setminus \{0\} \subset \{\lambda \in \mathbb{C} : \operatorname{Re} \lambda < -1\}.$$

In the same way we obtain the log-Sobolev inequality.

Theorem 4.5 *For any $\varphi \in C_b^1(\mathcal{H}^{-1})$ we have*

$$\int_{\mathcal{H}^{-1}} \varphi^2 \log \varphi^2 d\nu \leq \int_{\mathcal{H}^{-1}} |D_{\mathcal{H}^{-1}} \varphi(x)|_{\mathcal{H}^{-1}}^2 d\nu + \int_{\mathcal{H}^{-1}} \varphi^2 d\nu \log \left(\int_{\mathcal{H}^{-1}} \varphi^2 d\nu \right). \quad (4.15)$$

Remark 4.6 As already mentionned, all our results hold if we do not assume that the nonlinear term f in (4.1) is monotone but simply that it is the derivative of a semiconvex function, which is the case if f is a polynomial of odd degree with positive leading coefficient. In this case, in the proof of Theorem 4.1 we have to choose $\delta > 0$. The construction of the approximations U_α also has to be modified. The proof of theorem 4.3 and 4.5 do not use this assumption since it is known that ν_0 satisfy the spectral property and a log-Sobolev inequality also in that case.

References

- [1] **S. Albeverio and R. Høegh-Krohn**, *Dirichlet forms and diffusion processes on rigged Hilbert spaces*, Z. Wahrsch. verw. Geb., **40**, 1-57, 1977.
- [2] **S. Albeverio, V. Kondratiev and M. Röckner**, *Dirichlet operators via stochastic analysis*, Journal of Functional Analysis, **128**, n.1, 102–138, 1995.
- [3] **D. Bakry**, L'HYPERCONTRACTIVITÉ ET SON UTILISATION EN THÉORIE DES SEMI-GROUPES, Lectures on Probability Theory, LNM 1581, Springer, 1994.
- [4] **D. Blömker, S. Majer-Paape and T. Wanner**, *Spinodal Decomposition for the Cahn-Hilliard-Cook Equation*, Commun. Math. Phys., **223**, 553-582, 2001.
- [5] **A. Chojnowska-Michalik and B. Goldys**, *On regularity properties of nonsymmetric Ornstein-Uhlenbeck semigroup in L^p spaces*, Stoch. Stoch. Reports, **59**, 183-209, 1996.
- [6] **C. Cardon-Weber**, *Cahn-Hilliard stochastic equations: strict positivity of the density*, Stochastic and Stochastic Reports, **72**, 191-277, 2002.
- [7] **G. Da Prato & A. Debussche**, *Stochastic Cahn-Hilliard equation*, Nonlinear Anal. **26**, 2, 241-263, 1996.
- [8] **G. Da Prato, A. Debussche & B. Goldys**, *Invariant measures of non symmetric dissipative stochastic systems*, Probab. Theory Relat. Fields. **123**, 3, 355-380, 2000.
- [9] **G. Da Prato & L. Tubaro**, *Some results about dissipativity of Kolmogorov operators*, Czechoslovak Math. J., **51**, 4, 685-699, 2001.
- [10] **G. Da Prato & L. Tubaro**, *On a class of gradient systems with irregular potentials*, Infinite Dimensional Analysis, Quantum Probability and related topics, Vol. 4, 183–194, 2001.
- [11] **G. Da Prato & J. Zabczyk**, ERGODICITY FOR INFINITE DIMENSIONAL SYSTEMS, London Mathematical Society Lecture Notes, **229**, Cambridge University Press, 1996.
- [12] **G. Da Prato & J. Zabczyk**, SECOND ORDER PARTIAL DIFFERENTIAL EQUATIONS IN HILBERT SPACES, London Mathematical Society Lecture Notes, **293**, Cambridge University Press, 2002.
- [13] **A. Eberle**, UNIQUENESS AND NON-UNIQUENESS OF SINGULAR DIFFUSION OPERATORS, Lecture Notes in Mathematics, **1718**, Springer-Verlag, 1999.
- [14] **V. Liskevich and M. Röckner**, *Strong uniqueness for a class of infinite dimensional Dirichlet operators and application to stochastic quantization*, Ann. Scuola Norm. Sup. Pisa Cl. Sci., **27**, 69-91, 1999.
- [15] **G. Lumer & R. S. Phillips**, *Dissipative operators in a Banach space*, Pac. J. Math., **11**, 679-698, 1961.